

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **Goat anti Human IgA (Fc specific), conjugated with TRITC, Clone: [Polyclonal], Monoclonal NMB-GAHU/IGA(FC)/TRITC**

|                            |                                                                                                                                                                                                                                                                 |
|----------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Article Name               | Goat anti Human IgA (Fc specific), conjugated with TRITC, Clone:<br>[Polyclonal], Monoclonal                                                                                                                                                                    |
| Biozol Catalog Number      | NMB-GAHU/IGA(FC)/TRITC                                                                                                                                                                                                                                          |
| Supplier Catalog Number    | GAHu/IgA(Fc)/TRITC                                                                                                                                                                                                                                              |
| Alternative Catalog Number | NMB-GAHU/IGA(FC)/TRITC                                                                                                                                                                                                                                          |
| Manufacturer               | NordicMubio                                                                                                                                                                                                                                                     |
| Host                       | Goat                                                                                                                                                                                                                                                            |
| Category                   | Antikörper                                                                                                                                                                                                                                                      |
| Conjugation                | TRITC                                                                                                                                                                                                                                                           |
| Product Description        | The reactivity of the antiserum is directed to the Fc subunit of the IgA molecule which expresses strict isotypic (class) specificity. It does not react with any non-Ig protein in human serum, as tested by immunoelectrophoresis and double radial immu...   |
| Clonality                  | Monoclonal                                                                                                                                                                                                                                                      |
| Clone Designation          | [Polyclonal]                                                                                                                                                                                                                                                    |
| Buffer                     | TRITC-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity. It is reconstituted by adding 1 ml sterile distilled water, spun dow |

|                   |                                                                                                                                                                                            |
|-------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Source            | Purified polyclonal IgA and monoclonal IgA2 isolated from pooled human serum. Freund's complete adjuvant is used in the first step of the immunization procedure.                          |
| Form              | TRITC                                                                                                                                                                                      |
| Formula           | TRITC-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity. |
| Application Notes | ELISA, Immunocytochemistry, Immunohistochemistry (frozen), (In)direct immunofluorescence.                                                                                                  |