

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat anti Mouse IgD (Fc specific), Clone: [Polyclonal], Monoclonal NMB-GAM/IGD(FC)

| | |
|----------------------------|---|
| Article Name | Goat anti Mouse IgD (Fc specific), Clone: [Polyclonal], Monoclonal |
| Biozol Catalog Number | NMB-GAM/IGD(FC) |
| Supplier Catalog Number | GAM/IgD(Fc) |
| Alternative Catalog Number | NMB-GAM/IGD(FC) |
| Manufacturer | NordicMubio |
| Host | Goat |
| Category | Antikörper |
| Species Reactivity | Mouse |
| Conjugation | Unconjugated |
| Format | Antiserum |
| Target Specificity | IgD |
| Cross-Adsorption (MinX) | no cross-adsorbtion |
| Product Description | The reactivity of the antiserum is restricted to the Fc portion of the IgD molecule, which represents strict isotype (class specificity. In immunoelectrophoresis and radial immunodiffusion, using various antiserum concentrations against mouse serum b... |
| Clonality | Monoclonal |
| Clone Designation | [Polyclonal] |

| | |
|-------------------|--|
| Buffer | Delipidated, heat inactivated, lyophilized, stable whole antiserum No preservative added. Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal goat serum. No foreign proteins added. Reconstitute the lyophilized an |
| Source | Highly purified polyclonal and monoclonal IgD isolated from Balb/C and C57Bl serum. Freund's complete adjuvant is used in the first step of the immunization procedure. |
| Formula | Delipidated, heat inactivated, lyophilized, stable whole antiserum No preservative added. Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal goat serum. No foreign proteins added. |
| Antibody Type | Secondary Antibody |
| Application Notes | Precipitation assays. Dilutions may be prepared by adding phosphate buffered saline (PBS, pH 7.2). Repeated thawing and freezing should be avoided. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It will not affect the performance of the antiserum. Diluted antiserum should be stored at +4C, not refrozen, and preferably used the same day. |