

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **Goat anti Rabbit IgG IgA IgM (heavy and light chains), conjugated with TRITC, Clone: [Polyclonal], Monoclonal NMB-GAR/IG/TRITC**

Article Name	Goat anti Rabbit IgG IgA IgM (heavy and light chains), conjugated with TRITC, Clone: [Polyclonal], Monoclonal
Biozol Catalog Number	NMB-GAR/IG/TRITC
Supplier Catalog Number	GAR/Ig/TRITC
Alternative Catalog Number	NMB-GAR/IG/TRITC
Manufacturer	NordicMubio
Host	Goat
Category	Antikörper
Species Reactivity	Rabbit
Conjugation	TRITC
Format	IgG
Target Specificity	IgG+IgM+IgA (H+L)
Cross-Adsorption (MinX)	no cross-adsorbtion
Product Description	The reactivity of the antiserum is directed to the Fc and Fab subunits of the major immunoglobulins isotypes It does not react with any non-immunoglobulin in rabbit serum, as tested by immunoelectrophoresis and double radial immunodiffusion. To ident...
Clonality	Monoclonal
Clone Designation	[Polyclonal]

Buffer	TRITC-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity. It is reconstituted by adding 1 ml sterile distilled water, spun dow
Source	Purified immunoglobulin fractions containing IgG, IgA and IgM isolated from pooled rabbit serum. Freund's complete adjuvant is used in the first step of the immunization procedure.
Form	TRITC
Formula	TRITC-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity.
Antibody Type	Secondary Antibody
Application Notes	ELISA, Immunocytochemistry, Immunohistochemistry (frozen), (In)direct immunofluorescence.