

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### Goat anti Rat IgE (Fc specific), Clone: [Polyclonal], Monoclonal NMB-GARA/IGE(FC)

Article Name	Goat anti Rat IgE (Fc specific), Clone: [Polyclonal], Monoclonal
Biozol Catalog Number	NMB-GARA/IGE(FC)
Supplier Catalog Number	GARa/IgE(Fc)
Alternative Catalog Number	NMB-GARA/IGE(FC)
Manufacturer	NordicMubio
Host	Goat
Category	Antikörper
Species Reactivity	Rat
Conjugation	Unconjugated
Format	Antiserum
Target Specificity	IgE
Cross-Adsorption (MinX)	no cross-adsorbtion
Product Description	The reactivity of the antiserum is restricted to the Fc part of the IgE molecule, which represents strict isotype (class) specificity. In immunoelectrophoresis and radial immunodiffusion, using various antiserum concentrations against appropriate con...
Clonality	Monoclonal
Clone Designation	[Polyclonal]

Buffer	Delipidated, heat inactivated, lyophilized, stable whole antiserum No preservative added Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal goat serum. No foreign proteins added. Reconstitute the lyophilized ant
Source	Highly purified homogenous IgE isolated from rat serum. Freund's complete adjuvant is used in the first step of the immunization procedure.
Formula	Delipidated, heat inactivated, lyophilized, stable whole antiserum No preservative added Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal goat serum. No foreign proteins added.
Antibody Type	Secondary Antibody
Application Notes	Precipitation assays. In immunoelectrophoresis use 2 $\mu$ l serum or equivalent against 120 $\mu$ l antiserum. In double radial immunodiffusion (Ouchterlony) use a rosette arrangement with 10 $\mu$ l antiserum in 3 mm diameter center well and 2 $\mu$ l serum samples (neat and serially diluted) in 2 mm diameter peripheral wells.