

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### Goat anti Rat IgM (Fc specific), conjugated with TRITC, Clone: [Polyclonal], Monoclonal NMB-GARA/IGM(FC)/TRITC

Article Name	Goat anti Rat IgM (Fc specific), conjugated with TRITC, Clone: [Polyclonal], Monoclonal
Biozol Catalog Number	NMB-GARA/IGM(FC)/TRITC
Supplier Catalog Number	GARa/IgM(Fc)/TRITC
Alternative Catalog Number	NMB-GARA/IGM(FC)/TRITC
Manufacturer	NordicMubio
Host	Goat
Category	Antikörper
Species Reactivity	Rat
Conjugation	TRITC
Format	IgG
Target Specificity	IgM ( $\mu$ )
Cross-Adsorption (MinX)	no cross-adsorbtion
Product Description	The reactivity of the antiserum is directed to the Fc subunit of the IgM molecule which expresses strict isotypic (class) specificity. It does not react with purified IgG, IgG/Fab fragments and IgA or any non-Ig protein in rat serum, as tested by imm...
Clonality	Monoclonal
Clone Designation	[Polyclonal]

Buffer	TRITC-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity. It is reconstituted by adding 1 ml sterile distilled water, spun dow
Source	Purified pools of homogenous IgM isolated from rat serum. Freund's complete adjuvant is used in the first step of the immunization procedure.
Form	TRITC
Formula	TRITC-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity.
Antibody Type	Secondary Antibody
Application Notes	ELISA, Immunocytochemistry, Immunohistochemistry (frozen), (In)direct immunofluorescence.