

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat anti Turkey IgG (heavy and light chains), Clone: [Polyclonal], Monoclonal NMB-GATU/IGG(H+L)

Article Name	Goat anti Turkey IgG (heavy and light chains), Clone: [Polyclonal], Monoclonal
Biozol Catalog Number	NMB-GATU/IGG(H+L)
Supplier Catalog Number	GATu/IgG(H+L)
Alternative Catalog Number	NMB-GATU/IGG(H+L)
Manufacturer	NordicMubio
Host	Goat
Category	Antikörper
Species Reactivity	Aves
Conjugation	Unconjugated
Format	Antiserum
Target Specificity	IgG (H+L)
Cross-Adsorption (MinX)	no cross-adsorbtion
Product Description	In precipitating techniques as immunoelectrophoresis and double radial immunodiffusion to identify the presence of IgG, as secondary antibody to precipitate the immunoglobulin in normal turkey serum, to prepare an immunoadsorbent for the production o...
Clonality	Monoclonal
Clone Designation	[Polyclonal]

Buffer	Delipidated, heat inactivated lyophilized stable whole serum No preservative added, as it may interfere with the antibody activity. No foreign protein added. Total protein and IgG concentration in the antiserum are comparable to those of pooled goat serum
Source	Purified normal IgG isolated from pooled turkey serum. Freund's complete adjuvant is used in the first step of the immunization procedure.
Formula	Delipidated, heat inactivated lyophilized stable whole serum No preservative added, as it may interfere with the antibody activity. No foreign protein added. Total protein and IgG concentration in the antiserum are comparable to those of pooled goat serum
Antibody Type	Secondary Antibody
Application Notes	Precipitation assays. In immunoelectrophoresis use 2 μ l or equivalent against 120 μ l antiserum. In double radial immunodiffusion (Ouchterlony) use a rosette arrangement with 10 μ l antiserum in a 3 mm diameter centre well and 2 μ l serum samples (neat and diluted) in 2 mm diameter peripheral wells.