

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **Rabbit anti Duck IgG (heavy and light chains), Clone: [Polyclonal], Monoclonal NMB-RADU/IGG(H+L)**

Article Name	Rabbit anti Duck IgG (heavy and light chains), Clone: [Polyclonal], Monoclonal
Biozol Catalog Number	NMB-RADU/IGG(H+L)
Supplier Catalog Number	RADu/IgG(H+L)
Alternative Catalog Number	NMB-RADU/IGG(H+L)
Manufacturer	NordicMubio
Host	Rabbit
Category	Antikörper
Application	IP
Species Reactivity	Duck
Conjugation	Unconjugated
Format	Antiserum
Target Specificity	IgG (H+L)
Cross-Adsorption (MinX)	Bovine,Feline,Gallus,Canine,Goat,Equine,Human,Monkey,Mouse,Rat,S heep,Porcine
Product Description	The reactivity of the antiserum is directed to the Fc and Fab subunits of the IgG molecule. It includes a certain degree of reactivity with other immunoglobulins via the common Fab portion. It does not react with any non-Ig protein in duck serum, as ...
Clonality	Monoclonal

Clone Designation	[Polyclonal]
Buffer	Delipidated, heat inactivated, lyophilized, stable whole antiserum. No preservative added. Total protein and IgG concentration in the antiserum are comparable to those of pooled rabbit serum. No foreign proteins added. It is reconstituted by adding 1 ml s
Source	Purified normal IgG isolated from pooled duck serum. Freund's complete adjuvant is used in the first step of the immunization procedure.
Antibody Type	Secondary Antibody
Application Notes	Precipitation assays. In immunoelectrophoresis use 2 $\mu$ l or equivalent against 120 $\mu$ l antiserum. In double radial immunodiffusion (Ouchterlony) use a rosette arrangement with 10 $\mu$ l antiserum in a 3 mm diameter centre well and 2 $\mu$ l serum samples (neat and diluted) in 2 mm diameter peripheral wells. The optimum working dilution of the antibody for any specific application should be established by titration.