

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Rabbit anti Goat IgM (Fc specific), conjugated with Horseradish peroxidase, Clone: [Polyclonal], HRP, Monoclonal NMB-RAG/IGM(FC)/PO

| | |
|----------------------------|---|
| Article Name | Rabbit anti Goat IgM (Fc specific), conjugated with Horseradish peroxidase, Clone: [Polyclonal], HRP, Monoclonal |
| Biozol Catalog Number | NMB-RAG/IGM(FC)/PO |
| Supplier Catalog Number | RAG/IgM(Fc)/PO |
| Alternative Catalog Number | NMB-RAG/IGM(FC)/PO |
| Manufacturer | NordicMubio |
| Host | Rabbit |
| Category | Antikörper |
| Species Reactivity | Goat |
| Conjugation | HRP |
| Format | IgG |
| Target Specificity | IgM (μ) |
| Cross-Adsorption (MinX) | no cross-adsorbtion |
| Product Description | The reactivity of the antiserum is directed to the Fc subunit of the IgM molecule which expresses strict isotypic (class) specificity. It does not react with any non-Ig protein in goat serum, as tested by immunoelectrophoresis and double radial immun... |
| Clonality | Monoclonal |
| Clone Designation | [Polyclonal] |

| | |
|-------------------|---|
| Buffer | Peroxidase-coupled purified hyperimmune rabbit IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity. It is reconstituted by adding 1 ml sterile distilled water, s |
| Source | Purified normal IgM isolated from pooled goat serum. Freund's complete adjuvant is used in the first step of the immunization procedure. |
| Form | HRP |
| Formula | Peroxidase-coupled purified hyperimmune rabbit IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity. |
| Antibody Type | Secondary Antibody |
| Application Notes | ELISA, Immunocytochemistry, Immunohistochemistry (paraffin), Dot blot, Immunoblotting. |