

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Rabbit anti Goat IgM (heavy and light chains), Clone: [Polyclonal], Monoclonal NMB-RAG/IGM(H+L)

Article Name	Rabbit anti Goat IgM (heavy and light chains), Clone: [Polyclonal], Monoclonal
Biozol Catalog Number	NMB-RAG/IGM(H+L)
Supplier Catalog Number	RAG/IgM(H+L)
Alternative Catalog Number	NMB-RAG/IGM(H+L)
Manufacturer	NordicMubio
Host	Rabbit
Category	Antikörper
Species Reactivity	Goat
Conjugation	Unconjugated
Format	Antiserum
Target Specificity	IgM (H+L)
Cross-Adsorption (MinX)	no cross-adsorbtion
Product Description	The reactivity of the antiserum is directed to the Fc and Fab subunits of the IgM molecule. It includes a certain degree of reactivity with other immunoglobulins via the common Fab portion. It does not react with any non-Ig protein in goat serum, as ...
Clonality	Monoclonal
Clone Designation	[Polyclonal]

Buffer	Delipidated, heat inactivated lyophilized stable whole serum. No preservative added, as it may interfere with the antibody activity.No foreign protein added. Total protein and IgG concentration in the antiserum are comparable to those of pooled rabbit ser
Source	Purified normal IgM isolated from pooled goat serum. Freund's complete adjuvant is used in the first step of the immunization procedure.
Formula	Delipidated, heat inactivated lyophilized stable whole serum. No preservative added, as it may interfere with the antibody activity.No foreign protein added. Total protein and IgG concentration in the antiserum are comparable to those of pooled rabbit se
Antibody Type	Secondary Antibody
Application Notes	Precipitation assays. In immunoelectrophoresis use 2 μ l or equivalent against 120 μ l antiserum. In double radial immunodiffusion (Ouchterlony) use a rosette arrangement with 10 μ l antiserum in a 3 mm diameter centre well and 2 μ l serum samples (neat and diluted) in 2 mm diameter peripheral wells.