

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Rabbit anti Human beta thromboglobulin, Clone: [Polyclonal], Monoclonal NMB-RAHU/BTG

| | |
|----------------------------|---|
| Article Name | Rabbit anti Human beta thromboglobulin, Clone: [Polyclonal], Monoclonal |
| Biozol Catalog Number | NMB-RAHU/BTG |
| Supplier Catalog Number | RAHu/BTG |
| Alternative Catalog Number | NMB-RAHU/BTG |
| Manufacturer | NordicMubio |
| Host | Rabbit |
| Category | Antikörper |
| Product Description | The defined antibody specificity is restricted to beta-thromboglobulin as tested at the level of sensitivity of immunoprecipitation techniques. A single precipitin line is obtained in biddimensional electrophoresis against normal plasma and platelet ... |
| Clonality | Monoclonal |
| Clone Designation | [Polyclonal] |
| UniProt | P02775 |
| Buffer | Delipidated, heat inactivated lyophilized whole serum. Sodium azide 1 mg/ml. Total protein and IgG concentrations in the antiserum are comparable to those of normal pooled rabbit serum. No foreign proteins added. |
| Source | Beta-thromboglobulin is a platelet specific protein. In normal plasma mean concentrations of 30 ng/ml have been found. Highly purified beta-thromboglobulin has been isolated from platelets and is used for the immunization. Freund's complete adjuvant is used |

| | |
|-------------------|---|
| Formula | Delipidated, heat inactivated lyophilized whole serum. Sodium azide 1 mg/ml. Total protein and IgG concentrations in the antiserum are comparable to those of normal pooled rabbit serum. No foreign proteins added. |
| Application Notes | Precipitation assays. In electroimmunodiffusion the amount of antiserum required in the agarose gel is usually about 2% depending on the test arrangement. The amount of beta-thromboglobulin precipitated by 1 ml antiserum is approximately 0.2 mg. |