

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Rabbit anti Swine IgA (Fc specific), Clone: [Polyclonal], Monoclonal NMB-RASW/IGA(FC)

Article Name	Rabbit anti Swine IgA (Fc specific), Clone: [Polyclonal], Monoclonal
Biozol Catalog Number	NMB-RASW/IGA(FC)
Supplier Catalog Number	RASw/IgA(Fc)
Alternative Catalog Number	NMB-RASW/IGA(FC)
Manufacturer	NordicMubio
Host	Rabbit
Category	Antikörper
Species Reactivity	Porcine
Conjugation	Unconjugated
Format	Antiserum
Target Specificity	IgA
Cross-Adsorption (MinX)	no cross-adsorbtion
Product Description	The reactivity of the antiserum is restricted to the Fc part of the IgA molecule. In immunoelectrophoresis and radial immunodiffusion, using various antiserum concentrations against normal swine serum and milk, a single precipitin line is obtained wh...
Clonality	Monoclonal
Clone Designation	[Polyclonal]

Buffer	Delipidated, heat inactivated, lyophilized, stable whole antiserum. No preservative added Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal rabbit serum. No foreign proteins added. Reconstitute the lyophilized
Source	Highly purified normal IgA isolated from pooled swine serum. Freund's complete adjuvant is used in the first step of the immunization procedure.
Formula	Delipidated, heat inactivated, lyophilized, stable whole antiserum. No preservative added Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal rabbit serum. No foreign proteins added.
Antibody Type	Secondary Antibody
Application Notes	Precipitation assays. The lyophilized antiserum is shipped at ambient temperature and may be stored at +4C, prolonged storage at or below -20C. In immunoelectrophoresis use 2 µl serum or equivalent against 120 µl antiserum. In double radial immunodiffusion (Ouchterlony) use a rosette arrangement with 10 µl antiserum in 3 mm diameter center well and 2 µl serum samples (neat and serially diluted in 2 mm diameter peripheral wells).