

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Anti-Human IL-1alpha Antibody, Rabbit, Polyclonal ABT-ABG10235-U100

Article Name	Anti-Human IL-1alpha Antibody, Rabbit, Polyclonal
Biozol Catalog Number	ABT-ABG10235-U100
Supplier Catalog Number	ABG10235-U100
Alternative Catalog Number	ABT-ABG10235-U100-100UG
Manufacturer	Abcepta
Host	Rabbit
Category	Antikörper
Application	ELISA, IHC, WB
Species Reactivity	Human
Clonality	Polyclonal
Purity	Produced from sera of rabbits pre-immunized with highly pure (>98%) recombinant hIL-1alpha. Anti-Human IL-1alpha specific antibody was purified by affinity chromatography employing immobilized hIL-1alpha matrix.
Form	A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.
Antibody Type	Polyclonal Antibody

Application Notes	WesternBlot: To detect hIL-1alpha by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2 µg/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant hIL-1alpha is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.. Sandwich: To detect hIL-1alpha by sandwich ELISA (using 100 µl/well antibody solution) a concentration of 0.5 - 2.0 µg/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with BioGems Biotinylated Anti-Human IL-1alpha (60-001ABT) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant hIL-1alpha.. Immunohistochemistry: This antibody stained formalin-fixed, paraffin-embedded sections of normal human skin. The recommended concentration is 1.0 µg/ml overnight at 4C. An HRP-labeled polymer detection system was used with a non-alcohol soluble AEC chromogen. Enzyme induced antigen retrieval was performed with proteinase K. Optimal concentrations and conditions may vary.. Neutralization: To yield one-half maximal inhibition [ND50] of the biological activity of hIL-1alpha (20 ng/ml), a concentration of 1.0 - 2.0 µg/ml of this antibody is required.. Reconstitution: Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.
--------------------------	--