

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Anti-Human NNT-1/BCSF-3 Antibody, Rabbit, Polyclonal ABT-ABG10436-U100

| | |
|----------------------------|--|
| Article Name | Anti-Human NNT-1/BCSF-3 Antibody, Rabbit, Polyclonal |
| Biozol Catalog Number | ABT-ABG10436-U100 |
| Supplier Catalog Number | ABG10436-U100 |
| Alternative Catalog Number | ABT-ABG10436-U100-100UG |
| Manufacturer | Abcepta |
| Host | Rabbit |
| Category | Antikörper |
| Application | ELISA, WB |
| Species Reactivity | Human |
| Clonality | Polyclonal |
| Purity | Produced from sera of rabbits pre-immunized with highly pure (>98%) recombinant hNNT-1/BCSF-3. Anti-Human NNT-1/BCSF-3 specific antibody was purified by affinity chromatography employing immobilized hNNT-1/BCSF-3 matrix. |
| Form | A sterile filtered antibody solution was lyophilized from PBS, pH 7.2. |
| Antibody Type | Polyclonal Antibody |

Application Notes

WesternBlot: To detect hNNT-1/BCSF-3 by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2 µg/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant hNNT-1/BCSF-3 is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions. Sandwich: To detect hNNT-1/BCSF-3 by sandwich ELISA (using 100 µl/well antibody solution) a concentration of 0.5 - 2.0 µg/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with BioGems Biotinylated Anti-Human NNT-1/BCSF-3 (60-248BT) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant hNNT-1/BCSF-3.. Reconstitution: Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.