

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Anti-Human sCD40 Ligand Antibody, Goat, Polyclonal ABT-ABG10487-U050

Article Name	Anti-Human sCD40 Ligand Antibody, Goat, Polyclonal
Biozol Catalog Number	ABT-ABG10487-U050
Supplier Catalog Number	ABG10487-U050
Alternative Catalog Number	ABT-ABG10487-U050-50UG
Manufacturer	Abcepta
Host	Goat
Category	Antikörper
Application	ELISA, IHC, WB
Species Reactivity	Human
Clonality	Polyclonal
Purity	Produced from sera of goats pre-immunized with highly pure (>98%) recombinant hsCD40L. Anti-Human sCD40L specific antibody was purified by affinity chromatography employing immobilized hsCD40L matrix.
Form	A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.
Antibody Type	Polyclonal Antibody

Application Notes

WesternBlot: To detect hsCD40-Ligand by Western Blot analysis this antibody can be used at a concentration of 0.1- 0.2 µg/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant hsCD40-Ligand is 1.5-3.0 ng/lane, under either reducing or non-reducing conditions.. Sandwich: To detect hsCD40-Ligand by sandwich ELISA (using 100 µl/well antibody solution) a concentration of 0.5 - 2.0 µg/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with BioGems Biotinylated Anti-Human sCD40-Ligand (60-101) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant hsCD40-Ligand.. Immunohistochemistry: This antibody stained formalin-fixed, paraffin-embedded sections of human cervical squamous cell carcinoma. The recommended concentrations are 2.5 µg/ml - 5.0 µg/ml with an overnight incubation at 4C. An HRP-labeled polymer detection system was used with a DAB chromogen. The recommended antigen retrieval is heat induced antigen retrieval with a pH 6.0 Sodium Citrate buffer. Optimal concentrations and conditions may vary.. Reconstitution: Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.