

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Anti-Murine SDF-1alpha Antibody, Goat, Polyclonal ABT-ABG10501-U050

Article Name	Anti-Murine SDF-1alpha Antibody, Goat, Polyclonal
Biozol Catalog Number	ABT-ABG10501-U050
Supplier Catalog Number	ABG10501-U050
Alternative Catalog Number	ABT-ABG10501-U050-50UG
Manufacturer	Abcepta
Host	Goat
Category	Antikörper
Application	ELISA, WB
Species Reactivity	Mouse
Clonality	Polyclonal
Purity	Produced from sera of goats pre-immunized with highly pure recombinant Murine SDF-1alpha. Anti-Murine SDF-1alpha specific antibody was purified by affinity chromatography employing immobilized Murine SDF-1alpha matrix.
Form	A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.
Antibody Type	Polyclonal Antibody

Application Notes

WesternBlot: To detect Murine SDF-1alpha by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2 µg/ml. When used in conjunction with compatible secondary reagents, the detection limit for recombinant Murine SDF-1alpha is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions..

Sandwich: To detect Murine SDF-1alpha by sandwich ELISA (using 100 µl/well antibody solution) a concentration of 0.5 - 2.0 µg/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with BioGems Biotinylated Anti-Murine SDF-1alpha (61-120BT) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant Murine SDF-1alpha..

Neutralization: To yield one-half maximal inhibition [ND50] of the biological activity of Murine SDF-1alpha (100 ng/ml), a concentration of 2.5 - 3.5 µg/ml of this antibody is required..

Reconstitution: Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.