

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Catenin, beta (p120) Antibody, IgG1, Clone: [CTNNB1/1507], Mouse, Monoclonal NBT-1499-MSM7-P0

| | |
|----------------------------|---|
| Article Name | Catenin, beta (p120) Antibody, IgG1, Clone: [CTNNB1/1507], Mouse, Monoclonal |
| Biozol Catalog Number | NBT-1499-MSM7-P0 |
| Supplier Catalog Number | 1499-MSM7-P0 |
| Alternative Catalog Number | NBT-1499-MSM7-P0-20,NBT-1499-MSM7-P0-100 |
| Manufacturer | NeoBiotechnologies |
| Host | Mouse |
| Category | Antikörper |
| Application | IHC |
| Species Reactivity | Human, Mouse, Rat |
| Immunogen | Recombinant human beta-Catenin (p120) protein fragment |
| Product Description | Beta-catenin associates with the cytoplasmic portion of E-cadherin, which is necessary for the function of E-cadherin as an adhesion molecule. In normal tissues, beta-catenin is localized to the membrane of epithelial cells, consistent with its role ... |
| Clonality | Monoclonal |
| Clone Designation | [CTNNB1/1507] |
| Molecular Weight | 92kDa |
| Isotype | IgG1 |

| | |
|-------------------|---|
| NCBI | 1499 |
| UniProt | P35222 |
| Form | 200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Monoclonal Antibody |
| Application Notes | Immunohistochemistry (Formalin-fixed) (0.1-0.2ug/ml for 30 min at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes),Optimal dilution fo |