

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Glycophorin A / CD235a (Erythrocyte Marker) Antibody, IgG1, Clone: [GYPA/280], Mouse, Monoclonal NBT-2993-MSM4-P0

| | |
|----------------------------|---|
| Article Name | Glycophorin A / CD235a (Erythrocyte Marker) Antibody, IgG1, Clone: [GYPA/280], Mouse, Monoclonal |
| Biozol Catalog Number | NBT-2993-MSM4-P0 |
| Supplier Catalog Number | 2993-MSM4-P0 |
| Alternative Catalog Number | NBT-2993-MSM4-P0-20,NBT-2993-MSM4-P0-100 |
| Manufacturer | NeoBiotechnologies |
| Host | Mouse |
| Category | Antikörper |
| Application | IHC, WB |
| Species Reactivity | Human |
| Immunogen | Recombinant human full-length GYPA protein |
| Product Description | Recognizes a sialoglycoprotein of 39kDa, identified as glycophorin A (GPA). It is present on red blood cells (RBC) and erythroid precursor cells. It has been shown that glycophorin acts as the receptor for Sandei virus and parvovirus. Glycophorins A ... |
| Clonality | Monoclonal |
| Clone Designation | [GYPA/280] |
| Molecular Weight | 39kDa |
| Isotype | IgG1 |

| | |
|-------------------|---|
| NCBI | 2993 |
| UniProt | P02724 |
| Form | 200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Monoclonal Antibody |
| Application Notes | Western Blot (1-2ug/ml), Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 min |