

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Recombinant HSP60 (Heat Shock Protein 60) (Mitochondrial Marker) Antibody, Clone: [HSPD1/6498R], Rabbit, Monoclonal NBT-3329-RBM15-P1ABX

| | |
|----------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Article Name | Recombinant HSP60 (Heat Shock Protein 60) (Mitochondrial Marker) Antibody, Clone: [HSPD1/6498R], Rabbit, Monoclonal |
| Biozol Catalog Number | NBT-3329-RBM15-P1ABX |
| Supplier Catalog Number | 3329-RBM15-P1ABX |
| Alternative Catalog Number | NBT-3329-RBM15-P1ABX-100 |
| Manufacturer | NeoBiotechnologies |
| Host | Rabbit |
| Category | Antikörper |
| Application | IHC, WB |
| Species Reactivity | Bovine, Canine, Gallus, Hamster, Human, Monkey, Mouse, Porcine, Rabbit, Rat, Sheep |
| Immunogen | Recombinant human full-length HSP60 protein |
| Product Description | Recognizes a 60kDa protein, identified as the heat shock protein 60 (hsp60). Its epitope is localized between aa 383-447 of human hsp60. A wide variety of environmental and pathophysiological stressful conditions trigger the synthesis of a family of ... |
| Clonality | Monoclonal |
| Clone Designation | [HSPD1/6498R] |
| Molecular Weight | 60kDa |
| NCBI | 3329 |

| | |
|-------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| UniProt | P10809 |
| Form | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA at 1.0mg/ml. |
| Antibody Type | Recombinant Monoclonal Antibody |
| Application Notes | Western Blot (1-2ug/ml), Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95 °C followed by cooling at RT for |