

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Interferon gamma (IFNG) Antibody, IgG2b, Clone: [IFNG/466], Mouse, Monoclonal NBT-3458-MSM4-P0

| | |
|----------------------------|---|
| Article Name | Interferon gamma (IFNG) Antibody, IgG2b, Clone: [IFNG/466], Mouse, Monoclonal |
| Biozol Catalog Number | NBT-3458-MSM4-P0 |
| Supplier Catalog Number | 3458-MSM4-P0 |
| Alternative Catalog Number | NBT-3458-MSM4-P0-20,NBT-3458-MSM4-P0-100 |
| Manufacturer | NeoBiotechnologies |
| Host | Mouse |
| Category | Antikörper |
| Application | FC, IF, IHC |
| Species Reactivity | Human |
| Immunogen | Purified recombinant human full-length IFNG protein |
| Product Description | Interferon (IFN)- translated precursor is 166 amino acids, including the 23 amino acid secretory sequence. Multiple forms exist due to variable glycosylation and under non-denaturing conditions due to dimers and tetramers.... |
| Clonality | Monoclonal |
| Clone Designation | [IFNG/466] |
| Molecular Weight | 20-25kDa |
| Isotype | IgG2b |

| | |
|-------------------|---|
| NCBI | 3458 |
| UniProt | P01579 |
| Form | 200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Monoclonal Antibody |
| Application Notes | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 min at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes)Optimal dilution for |