

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Recombinant TACSTD2 / TROP2 (Epithelial Marker) Antibody, IgG1, Clone: [rTACSTD2/6397], Mouse, Monoclonal NBT-4070-MSM7-P0

| | |
|----------------------------|---|
| Article Name | Recombinant TACSTD2 / TROP2 (Epithelial Marker) Antibody, IgG1, Clone: [rTACSTD2/6397], Mouse, Monoclonal |
| Biozol Catalog Number | NBT-4070-MSM7-P0 |
| Supplier Catalog Number | 4070-MSM7-P0 |
| Alternative Catalog Number | NBT-4070-MSM7-P0-20,NBT-4070-MSM7-P0-100 |
| Manufacturer | NeoBiotechnologies |
| Host | Mouse |
| Category | Antikörper |
| Application | IHC, WB |
| Species Reactivity | Guinea pig, Hamster, Human, Mouse, Rabbit, Rat |
| Immunogen | Recombinant fragment of human TACSTD2 protein (around aa 31-274) (exact sequence is proprietary) |
| Product Description | TACSTD2 is a cell surface glycoprotein receptor. It is a single pass type I membrane protein containing one thyroglobulin type-1 domain, an epidermal growth factor-like repeat, a phosphatidylinositol binding site and tyrosine phosphorylation sites ne... |
| Clonality | Monoclonal |
| Clone Designation | [rTACSTD2/6397] |
| Molecular Weight | 40kDa |

| | |
|-------------------|---|
| Isotype | IgG1 |
| NCBI | 4070 |
| UniProt | P09758 |
| Form | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Recombinant Monoclonal Antibody |
| Application Notes | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95 °C followed by cooling at RT for 20 minutes),Optimal diluti |