

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **NKX2.2 (Neuroendocrine & Ewings Sarcoma Marker) Antibody, Rabbit, Polyclonal** **NBT-4821-RBP5-P0**

|                            |                                                                                                                                                                                                                                                               |
|----------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Article Name               | NKX2.2 (Neuroendocrine & Ewings Sarcoma Marker) Antibody, Rabbit, Polyclonal                                                                                                                                                                                  |
| Biozol Catalog Number      | NBT-4821-RBP5-P0                                                                                                                                                                                                                                              |
| Supplier Catalog Number    | 4821-RBP5-P0                                                                                                                                                                                                                                                  |
| Alternative Catalog Number | NBT-4821-RBP5-P0-20,NBT-4821-RBP5-P0-100                                                                                                                                                                                                                      |
| Manufacturer               | NeoBiotechnologies                                                                                                                                                                                                                                            |
| Host                       | Rabbit                                                                                                                                                                                                                                                        |
| Category                   | Antikörper                                                                                                                                                                                                                                                    |
| Application                | IHC                                                                                                                                                                                                                                                           |
| Species Reactivity         | Human                                                                                                                                                                                                                                                         |
| Immunogen                  | Recombinant human NKX2.2 protein fragment                                                                                                                                                                                                                     |
| Product Description        | Expression of NKX2.2 has been found in neuroendocrine tumors of the gut, making it a potential marker for the study of gastrointestinal neuroendocrine tumors. More recently, NKX2.2 protein was identified as a target of EWS-FLI-1, the fusion protein s... |
| Clonality                  | Polyclonal                                                                                                                                                                                                                                                    |
| Molecular Weight           | 40-50kDa                                                                                                                                                                                                                                                      |
| NCBI                       | <a href="#">4821</a>                                                                                                                                                                                                                                          |
| UniProt                    | <a href="#">O95096</a>                                                                                                                                                                                                                                        |

|                   |                                                                                                                                                                                                                                                               |
|-------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Form              | 200ug/ml of Ab Purified by Protein A. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.                                                                                                                      |
| Application Notes | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes), Optimal dilution f |