

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

GTF2H2C (Transcription Factor) Antibody, IgG2c, Clone: [PCRP-GTF2H2C-2C9], Mouse, Monoclonal NBT-730394-MSM1-P1ABX

| | |
|----------------------------|--|
| Article Name | GTF2H2C (Transcription Factor) Antibody, IgG2c, Clone: [PCRP-GTF2H2C-2C9], Mouse, Monoclonal |
| Biozol Catalog Number | NBT-730394-MSM1-P1ABX |
| Supplier Catalog Number | 730394-MSM1-P1ABX |
| Alternative Catalog Number | NBT-730394-MSM1-P1ABX-100 |
| Manufacturer | NeoBiotechnologies |
| Host | Mouse |
| Category | Antikörper |
| Application | FC, IF, IHC, WB |
| Species Reactivity | Human |
| Immunogen | Recombinant full-length human GTF2H2C protein |
| Product Description | Initiation of transcription from protein-coding genes in eukaryotes is a complex process that requires RNA polymerase II, as well as families of basal transcription factors. Binding of the factor TFIID (TBP) to the TATA box is believed to be the first... |
| Clonality | Monoclonal |
| Clone Designation | [PCRP-GTF2H2C-2C9] |
| Molecular Weight | 44kDa |
| Isotype | IgG2c |

| | |
|-------------------|---|
| NCBI | 730394 |
| UniProt | Q6P1K8 |
| Form | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Monoclonal Antibody |
| Application Notes | Flow Cytometry (1-2ug/million cells), Immunofluorescence (1-2ug/ml), Western Blot (1-2ug/ml), Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1m |