

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Histone Deacetylase 1 (HDAC3) Antibody, IgG2b, Clone: [PCRP-HDAC3-3C9], Mouse, Monoclonal NBT-8841-MSM2-P0

| | |
|----------------------------|---|
| Article Name | Histone Deacetylase 1 (HDAC3) Antibody, IgG2b, Clone: [PCRP-HDAC3-3C9], Mouse, Monoclonal |
| Biozol Catalog Number | NBT-8841-MSM2-P0 |
| Supplier Catalog Number | 8841-MSM2-P0 |
| Alternative Catalog Number | NBT-8841-MSM2-P0-20,NBT-8841-MSM2-P0-100 |
| Manufacturer | NeoBiotechnologies |
| Host | Mouse |
| Category | Antikörper |
| Application | ELISA, FC, IF, WB |
| Species Reactivity | Human |
| Immunogen | Recombinant full-length human HDAC3 protein |
| Product Description | In the intact cell, DNA closely associates with histones and other nuclear proteins to form chromatin. The remodeling of chromatin is believed to be a critical component of transcriptional regulation and a major source of this remodeling is brought a... |
| Clonality | Monoclonal |
| Clone Designation | [PCRP-HDAC3-3C9] |
| Molecular Weight | 49kDa |
| Isotype | IgG2b |

| | |
|-------------------|--|
| NCBI | 8841 |
| UniProt | O15379 |
| Form | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Monoclonal Antibody |
| Application Notes | ELISA (For coating, order antibody without BSA), Western Blot (1-2ug/ml), Immunoprecipitation (1-2ug per 100-500ug of total protein (1ml of cell lysate)), Flow Cytometry (1-2ug/million cells), Immunofluorescence (1-2ug/ml), Optimal dilution for a speci |