

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Recombinant Napsin A (Lung Adenocarcinoma Marker) Antibody, Clone: [NAPSA/7043R], Rabbit, Monoclonal NBT-9476-RBM17-P1ABX

| | |
|----------------------------|--|
| Article Name | Recombinant Napsin A (Lung Adenocarcinoma Marker) Antibody, Clone: [NAPSA/7043R], Rabbit, Monoclonal |
| Biozol Catalog Number | NBT-9476-RBM17-P1ABX |
| Supplier Catalog Number | 9476-RBM17-P1ABX |
| Alternative Catalog Number | NBT-9476-RBM17-P1ABX-100 |
| Manufacturer | NeoBiotechnologies |
| Host | Rabbit |
| Category | Antikörper |
| Application | IHC, WB |
| Species Reactivity | Human |
| Immunogen | Recombinant human Napsin-A protein fragment (aa1-100) (exact sequence is proprietary) |
| Product Description | Napsin is a pepsin-like aspartic proteinase connected with maturation of surfactant protein B. There are two closely related napsins, napsin A and napsin B. Napsin A is expressed as a single chain protein. Immunohistochemical studies revealed high exp... |
| Clonality | Monoclonal |
| Clone Designation | [NAPSA/7043R] |
| Molecular Weight | 37kDa |
| NCBI | 9476 |

| | |
|-------------------|---|
| UniProt | O96009 |
| Form | 200ug/ml of Ab purified from rabbit anti-serum by Protein A. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA at 1.0mg/ml. |
| Antibody Type | Recombinant Monoclonal Antibody |
| Application Notes | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95 °C followed by cooling at RT for 20 minutes),Optimal diluti |