

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

CD43 (T-Cell Marker) (rSPN/839), CF640R conjugate, 0.1mg/mL, IgG1, Clone: [rSPN/839], Mouse, Monoclonal BOT-BNC402220-500

| | |
|----------------------------|---|
| Article Name | CD43 (T-Cell Marker) (rSPN/839), CF640R conjugate, 0.1mg/mL, IgG1, Clone: [rSPN/839], Mouse, Monoclonal |
| Biozol Catalog Number | BOT-BNC402220-500 |
| Supplier Catalog Number | BNC402220-500 |
| Alternative Catalog Number | BOT-BNC402220-500-500UL |
| Manufacturer | Biotium |
| Host | Mouse |
| Category | Antikörper |
| Application | FC, IF, IHC, WB |
| Species Reactivity | Human |
| Immunogen | Recombinant full-length human CD43 protein |
| Conjugation | CF640R |
| Product Description | This antibody recognizes a cell surface glycoprotein of 95/115/135 kDa (depending upon the extent of glycosylation), identified as CD43 . Epitope of MAb Bra7G is clearly different from that of MAb rSPN/839, called b as opposed to a for rSPN/839. 70-9... |
| Clonality | Monoclonal |
| Concentration | 0.1 mg/mL |
| Clone Designation | [rSPN/839] |

| | |
|-------------------|---|
| Molecular Weight | 95, 115, or 135 kDa |
| Isotype | IgG1 |
| UniProt | P16150 |
| Buffer | PBS, 0.1% BSA, 0.05% azide |
| Source | Animal |
| Application Notes | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 0.5-1 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user |