

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Moesin (rMSN/492), CF640R conjugate, 0.1mg/mL, Clone: [rMSN/492], Mouse, Monoclonal BOT-BNC402307-100

| | |
|----------------------------|---|
| Article Name | Moesin (rMSN/492), CF640R conjugate, 0.1mg/mL, Clone: [rMSN/492], Mouse, Monoclonal |
| Biozol Catalog Number | BOT-BNC402307-100 |
| Supplier Catalog Number | BNC402307-100 |
| Alternative Catalog Number | BOT-BNC402307-100-100UL |
| Manufacturer | Biotium |
| Host | Mouse |
| Category | Antikörper |
| Application | IF, IHC, WB |
| Species Reactivity | Human, Rat |
| Immunogen | Recombinant full-length human Moesin protein |
| Conjugation | CF640R |
| Product Description | This antibody recognizes 78 kDa moesin protein. Moesin, a member of the talin-4.1 superfamily, is a linking protein of the sub-membranous actin cytoskeleton. It is expressed in variable amounts in cells of different phenotypes such as macrophages, ly... |
| Clonality | Monoclonal |
| Concentration | 0.1 mg/mL |
| Clone Designation | [rMSN/492] |

| | |
|-------------------|--|
| Molecular Weight | 78 kDa |
| UniProt | P26038 |
| Buffer | PBS, 0.1% BSA, 0.05% azide |
| Source | Animal |
| Application Notes | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 0.5-1 ug/mL for 30 minutes at RT Western Blot 0.5-2 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user |