

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

CD81 / TAPA-1 (C81/2885R), CF647 conjugate, 0.1mg/mL, Clone: [C81/2885R], Rabbit, Monoclonal BOT-BNC472885-100

| | |
|----------------------------|---|
| Article Name | CD81 / TAPA-1 (C81/2885R), CF647 conjugate, 0.1mg/mL, Clone: [C81/2885R], Rabbit, Monoclonal |
| Biozol Catalog Number | BOT-BNC472885-100 |
| Supplier Catalog Number | BNC472885-100 |
| Alternative Catalog Number | BOT-BNC472885-100-100UL |
| Manufacturer | Biotium |
| Host | Rabbit |
| Category | Antikörper |
| Application | Exosome Staining, FC, IHC |
| Species Reactivity | Human, Mouse, Rat |
| Immunogen | B-Cell line derived from a Burkitt lymphoma |
| Conjugation | CF647 |
| Product Description | This antibody recognizes a protein of 26 kDa, identified as CD81 (Workshop VI, Code CD81.1). CD81 has a very broad cellular distribution, being expressed on T- and B-lymphocytes, NK cells, thymocytes, eosinophils, fibroblasts, epithelial and endothel... |
| Clonality | Monoclonal |
| Concentration | 0.1 mg/mL |
| Clone Designation | [C81/2885R] |

| | |
|-------------------|--|
| Molecular Weight | 26 kDa |
| UniProt | P60033 |
| Buffer | PBS, 0.1% BSA, 0.05% azide |
| Source | Animal |
| Application Notes | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 2-4 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris with 1 mM EDTA pH 9.0 for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user |