

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **Uroplakin 1A (Urothelial Differentiation Marker)(UPK1A/2924), CF647 conjugate, 0.1mg/mL, Clone: [UPK1A/2924], Mouse, Monoclonal BOT-BNC472924-500**

Article Name	Uroplakin 1A (Urothelial Differentiation Marker)(UPK1A/2924), CF647 conjugate, 0.1mg/mL, Clone: [UPK1A/2924], Mouse, Monoclonal
Biozol Catalog Number	BOT-BNC472924-500
Supplier Catalog Number	BNC472924-500
Alternative Catalog Number	BOT-BNC472924-500-500UL
Manufacturer	Biotium
Host	Mouse
Category	Antikörper
Species Reactivity	Human
Immunogen	Recombinant fragment (around aa 114-173) of human Uroplakin 1A (UPK1A) protein (exact sequence is proprietary)
Conjugation	CF647
Product Description	The protein encoded by this gene is a member of the transmembrane 4 superfamily, also known as the tetraspanin family. Most of these members are cell-surface proteins that are characterized by the presence of four hydrophobic domains. The proteins me...
Clonality	Monoclonal
Concentration	0.1 mg/mL
Clone Designation	[UPK1A/2924]

Molecular Weight	29 kDa
UniProt	<a href="#">O00322</a>
Buffer	PBS, 0.1% BSA, 0.05% azide
Source	Animal
Application Notes	For coating for ELISA, order Ab without BSA Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Optimal dilution and staining procedure for a specific application should be determined by user Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry