

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Neurofilament (NF-L) (Neuronal Marker) (NEFL/2983R), CF647 conjugate, 0.1mg/mL, Clone: [NEFL/2983R], Rabbit, Monoclonal BOT-BNC472983-100

| | |
|----------------------------|---|
| Article Name | Neurofilament (NF-L) (Neuronal Marker) (NEFL/2983R), CF647 conjugate, 0.1mg/mL, Clone: [NEFL/2983R], Rabbit, Monoclonal |
| Biozol Catalog Number | BOT-BNC472983-100 |
| Supplier Catalog Number | BNC472983-100 |
| Alternative Catalog Number | BOT-BNC472983-100-100UL |
| Manufacturer | Biotium |
| Host | Rabbit |
| Category | Antikörper |
| Application | IHC |
| Species Reactivity | Bovine, Gallus, Human, Porcine, Rat |
| Immunogen | Recombinant full-length human NEFL protein |
| Conjugation | CF647 |
| Product Description | This MAb reacts with a 68 kDa protein, identified as light sub-unit of neurofilaments (NF-L). Neurofilaments make up the main structural elements of axons and dendrites and are found in neurons, peripheral nerves, and sympathetic ganglion cells. Neur... |
| Clonality | Monoclonal |
| Concentration | 0.1 mg/mL |
| Clone Designation | [NEFL/2983R] |

| | |
|-------------------|--|
| Molecular Weight | 68 kDa |
| UniProt | P07196 |
| Buffer | PBS, 0.1% BSA, 0.05% azide |
| Source | Animal |
| Application Notes | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 0.25-0.5 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user |