

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **Cytokeratin, pan (Cocktail PAN-CK), 0.2mg/mL, Clone: [Cocktail PAN-CK], Mouse, Monoclonal BOT-BNUB0999-100**

|                            |                                                                                                                                                                                                                                                               |
|----------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Article Name               | Cytokeratin, pan (Cocktail PAN-CK), 0.2mg/mL, Clone: [Cocktail PAN-CK], Mouse, Monoclonal                                                                                                                                                                     |
| Biozol Catalog Number      | BOT-BNUB0999-100                                                                                                                                                                                                                                              |
| Supplier Catalog Number    | BNUB0999-100                                                                                                                                                                                                                                                  |
| Alternative Catalog Number | BOT-BNUB0999-100-100UL                                                                                                                                                                                                                                        |
| Manufacturer               | Biotium                                                                                                                                                                                                                                                       |
| Host                       | Mouse                                                                                                                                                                                                                                                         |
| Category                   | Antikörper                                                                                                                                                                                                                                                    |
| Application                | FC, IF, IHC                                                                                                                                                                                                                                                   |
| Species Reactivity         | Bovine, Canine, Gallus, Human, Monkey, Mouse, Rabbit, Rat                                                                                                                                                                                                     |
| Immunogen                  | Human epidermal keratin                                                                                                                                                                                                                                       |
| Product Description        | Twenty human keratins are resolved with two-dimensional gel electrophoresis into acidic (pI 6.0) subfamilies. This antibody cocktail recognizes acidic (Type I or LMW) and basic (Type II or HMW) cytokeratins, with 67 kDa (CK1), 64 kDa (CK3), 59 kDa (C... |
| Clonality                  | Monoclonal                                                                                                                                                                                                                                                    |
| Concentration              | 0.2 mg/mL                                                                                                                                                                                                                                                     |
| Clone Designation          | [Cocktail PAN-CK]                                                                                                                                                                                                                                             |
| Molecular Weight           | 40-67 kDa (Multiple)                                                                                                                                                                                                                                          |

|                   |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                |
|-------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| UniProt           | <a href="#">Q7Z794</a>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |
| Buffer            | PBS, 0.05% BSA, 0.05% azide                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    |
| Source            | Animal                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |
| Application Notes | <p>Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user</p> |