

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

CD44v4 (Marker of Tumor Metastasis) (CD44v4/1700R), 0.2mg/mL, Clone: [CD44v4/1700R], Rabbit, Monoclonal BOT-BNUB1700-500

Article Name	CD44v4 (Marker of Tumor Metastasis) (CD44v4/1700R), 0.2mg/mL, Clone: [CD44v4/1700R], Rabbit, Monoclonal
Biozol Catalog Number	BOT-BNUB1700-500
Supplier Catalog Number	BNUB1700-500
Alternative Catalog Number	BOT-BNUB1700-500-500UL
Manufacturer	Biotium
Host	Rabbit
Category	Antikörper
Application	IHC
Species Reactivity	Human
Immunogen	Recombinant fragment corresponding to the v4 domain of human CD44 (exact sequence is proprietary)
Product Description	This antibody recognizes an epitope encoded by exon v4 on the variant portion of human CD44. The CD44 molecule belongs to a family of cellular adhesion molecules found on a wide range of normal and malignant cells in epithelial, mesothelial and hemop...
Clonality	Monoclonal
Concentration	0.2 mg/mL
Clone Designation	[CD44v4/1700R]
Molecular Weight	80-95 kDa

UniProt	P16070
Buffer	PBS, 0.05% BSA, 0.05% azide
Source	Animal
Application Notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin): 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user