

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **TACSTD2 / TROP2 (Epithelial Marker) (TACSTD2/2152), 0.2mg/mL, Clone: [TACSTD2/2152], Mouse, Monoclonal BOT-BNUB2152-500**

|                            |                                                                                                                                                                                                                                                               |
|----------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Article Name               | TACSTD2 / TROP2 (Epithelial Marker) (TACSTD2/2152), 0.2mg/mL, Clone: [TACSTD2/2152], Mouse, Monoclonal                                                                                                                                                        |
| Biozol Catalog Number      | BOT-BNUB2152-500                                                                                                                                                                                                                                              |
| Supplier Catalog Number    | BNUB2152-500                                                                                                                                                                                                                                                  |
| Alternative Catalog Number | BOT-BNUB2152-500-500UL                                                                                                                                                                                                                                        |
| Manufacturer               | Biotium                                                                                                                                                                                                                                                       |
| Host                       | Mouse                                                                                                                                                                                                                                                         |
| Category                   | Antikörper                                                                                                                                                                                                                                                    |
| Application                | IF, IHC                                                                                                                                                                                                                                                       |
| Species Reactivity         | Human                                                                                                                                                                                                                                                         |
| Immunogen                  | Recombinant fragment of human TACSTD2 protein (around aa 31-274) (exact sequence is proprietary)                                                                                                                                                              |
| Product Description        | TACSTD2 is a cell surface glycoprotein receptor. It is a single pass type I membrane protein containing one thyroglobulin type-1 domain, an epidermal growth factor-like repeat, a phosphatidylinositol binding site and tyrosine phosphorylation sites ne... |
| Clonality                  | Monoclonal                                                                                                                                                                                                                                                    |
| Concentration              | 0.2 mg/mL                                                                                                                                                                                                                                                     |
| Clone Designation          | [TACSTD2/2152]                                                                                                                                                                                                                                                |

|                   |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |
|-------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Molecular Weight  | 40 kDa                                                                                                                                                                                                                                                                                                                                                                                                                                                                           |
| UniProt           | <a href="#">P09758</a>                                                                                                                                                                                                                                                                                                                                                                                                                                                           |
| Buffer            | PBS, 0.05% BSA, 0.05% azide                                                                                                                                                                                                                                                                                                                                                                                                                                                      |
| Source            | Animal                                                                                                                                                                                                                                                                                                                                                                                                                                                                           |
| Application Notes | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user |