

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **CD15 / FUT4 / Lewis x(FUT4/815 + BRA-4F1), 1mg/mL, IgM, Clone: [FUT4/815 BRA-4F1], Mouse, Monoclonal BOT-BNUM0562-50**

Article Name	CD15 / FUT4 / Lewis x(FUT4/815 + BRA-4F1), 1mg/mL, IgM, Clone: [FUT4/815 BRA-4F1], Mouse, Monoclonal
Biozol Catalog Number	BOT-BNUM0562-50
Supplier Catalog Number	BNUM0562-50
Alternative Catalog Number	BOT-BNUM0562-50-50UL
Manufacturer	Biotium
Host	Mouse
Category	Antikörper
Application	IHC
Species Reactivity	Human
Immunogen	Purified human neutrophils (FUT4/815), Myelomonocytic leukemia cells (BRA-4F1)
Product Description	CD15 plays a role in mediating phagocytosis, bactericidal activity, and chemotaxis. It is present on >95% of granulocytes including neutrophils and eosinophils and to a lesser degree on monocytes. In addition, CD15 is expressed in Reed-Sternberg cell...
Clonality	Monoclonal
Concentration	1 mg/mL
Clone Designation	[FUT4/815 BRA-4F1]
Molecular Weight	~220 kDa (Glycoprotein)

Isotype	IgM
UniProt	<a href="#">P22083</a>
Buffer	PBS, no BSA, no azide
Source	Animal
Application Notes	Immunohistology (formalin) Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris with 1 mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 minutes Immunofluorescence 0.5-1 ug/mL Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user