

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Hepatocyte Specific(OCH1E5), 1mg/mL, IgG1, Clone: [OCH1E5], Mouse, Monoclonal BOT-BNUM0671-50

| | |
|----------------------------|---|
| Article Name | Hepatocyte Specific(OCH1E5), 1mg/mL, IgG1, Clone: [OCH1E5], Mouse, Monoclonal |
| Biozol Catalog Number | BOT-BNUM0671-50 |
| Supplier Catalog Number | BNUM0671-50 |
| Alternative Catalog Number | BOT-BNUM0671-50-50UL |
| Manufacturer | Biotium |
| Host | Mouse |
| Category | Antikörper |
| Application | IHC |
| Species Reactivity | Canine, Human |
| Immunogen | Extract of a formalin-fixed, rejected-allograft of a human liver |
| Product Description | Hepatocyte Specific Antigen, also called Hepatocyte Paraffin 1 or HepPar1, localizes to the mitochondria of hepatocytes. It is a sensitive marker for distinguishing hepatocellular carcinomas (HCC) from other metastatic carcinomas as well as cholangio... |
| Clonality | Monoclonal |
| Concentration | 1 mg/mL |
| Clone Designation | [OCH1E5] |
| Molecular Weight | Not Known |

| | |
|-------------------|---|
| Isotype | IgG1 |
| UniProt | Not Known |
| Buffer | PBS, no BSA, no azide |
| Source | Animal |
| Application Notes | Immunohistology formalin-fixed 1-2 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris with 1 mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 minutes Immunofluorescence 0.5-1.0 ug/mL Optimal dilution for a specific application should be determined by user |