

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

c-Myc Oncoprotein(MYC275 + MYC909), 1mg/mL, Clone: [MYC275 MYC909], Mouse, Monoclonal BOT-BNUM1269-50

Article Name	c-Myc Oncoprotein(MYC275 + MYC909), 1mg/mL, Clone: [MYC275 MYC909], Mouse, Monoclonal
Biozol Catalog Number	BOT-BNUM1269-50
Supplier Catalog Number	BNUM1269-50
Alternative Catalog Number	BOT-BNUM1269-50-50UL
Manufacturer	Biotium
Host	Mouse
Category	Antikörper
Application	IHC
Species Reactivity	Human
Immunogen	Recombinant human c-myc protein
Product Description	This antibody recognizes a transcription factor of 64-67 kDa, identified as c-myc. This MAb shows no cross-reaction with v-myc. c-myc is involved in the control of cell proliferation and differentiation and is amplified and/or over-expressed in a var...
Clonality	Monoclonal
Concentration	1 mg/mL
Clone Designation	[MYC275 MYC909]
Molecular Weight	62-64 kDa

UniProt	P01106
Buffer	PBS, no BSA, no azide
Source	Animal
Application Notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistochemistry (formalin-fixed): 1-2 ug/mL for 30 minutes at RT Flow cytometry: 0.5-1 ug/million cells Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris with 1 mM EDTA pH 9.0 for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user