

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **CD56 / NCAM1 / NKH1 (Neuronal Cell Marker) (NCAM1/1496), 1mg/mL, Clone: [NCAM1/1496], Mouse, Monoclonal BOT-BNUM1496-50**

Article Name	CD56 / NCAM1 / NKH1 (Neuronal Cell Marker) (NCAM1/1496), 1mg/mL, Clone: [NCAM1/1496], Mouse, Monoclonal
Biozol Catalog Number	BOT-BNUM1496-50
Supplier Catalog Number	BNUM1496-50
Alternative Catalog Number	BOT-BNUM1496-50-50UL
Manufacturer	Biotium
Host	Mouse
Category	Antikörper
Species Reactivity	Human
Immunogen	Recombinant human NCAM1 protein
Product Description	This MAb reacts with an extracellular domain (close to transmembrane) of CD56/NCAM. Three isoforms of neural cell adhesion molecule (NCAM) are produced by differential splicing of the RNA transcript from a single gene. The 135 kDa isoform is the basi...
Clonality	Monoclonal
Concentration	1 mg/mL
Clone Designation	[NCAM1/1496]
Molecular Weight	125 kDa 145 180
UniProt	<a href="#">P13591</a>

Buffer	PBS, no BSA, no azide
Source	Animal
Application Notes	For coating for ELISA, order Ab without BSA Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Optimal dilution and staining procedure for a specific application should be determined by user Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry