

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

CA19-9/Sialyl Lewisa (GI Tumor Marker) (CA19.9/1390R), CF740 conjugate, 0.1mg/mL, Clone: [CA19.9/1390R], CF 740, Rabbit, Monoclonal BOT-BNC741390-500

| | |
|----------------------------|---|
| Article Name | CA19-9/Sialyl Lewisa (GI Tumor Marker) (CA19.9/1390R), CF740 conjugate, 0.1mg/mL, Clone: [CA19.9/1390R], CF 740, Rabbit, Monoclonal |
| Biozol Catalog Number | BOT-BNC741390-500 |
| Supplier Catalog Number | BNC741390-500 |
| Alternative Catalog Number | BOT-BNC741390-500-500UL |
| Manufacturer | Biotium |
| Host | Rabbit |
| Category | Antikörper |
| Application | IHC |
| Species Reactivity | Human |
| Immunogen | Purified human CA19-9 protein |
| Conjugation | CF 740 |
| Product Description | CA19-9, a carbohydrate epitope expressed on a high MW (>400 kDa) mucin glycoprotein, is a sialyl Lewisa structure which is synthesized from type 1 blood group precursor chains and is present in individuals expressing the Lewisa and/or Lewisb blood gr... |
| Clonality | Monoclonal |
| Concentration | 0.1 mg/mL |
| Clone Designation | [CA19.9/1390R] |

| | |
|-------------------|---|
| Molecular Weight | >400 kDa |
| UniProt | Not Known |
| Buffer | PBS, 0.1% rBSA, 0.05% azide |
| Source | Animal |
| Application Notes | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence 0.5-1 ug/mL Immunohistology (formalin) 5-10 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |