

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

CD31 / PECAM-1 (Endothelial Cell Marker) (C31/1395R), CF740 conjugate, 0.1mg/mL, Clone: [C31/1395R], CF 740, Rabbit, Monoclonal BOT-BNC741395-100

| | |
|----------------------------|---|
| Article Name | CD31 / PECAM-1 (Endothelial Cell Marker) (C31/1395R), CF740 conjugate, 0.1mg/mL, Clone: [C31/1395R], CF 740, Rabbit, Monoclonal |
| Biozol Catalog Number | BOT-BNC741395-100 |
| Supplier Catalog Number | BNC741395-100 |
| Alternative Catalog Number | BOT-BNC741395-100-100UL |
| Manufacturer | Biotium |
| Host | Rabbit |
| Category | Antikörper |
| Application | FC, IHC, WB |
| Species Reactivity | Human |
| Immunogen | Human recombinant full-length CD31 protein |
| Conjugation | CF 740 |
| Product Description | CD31 (PECAM-1) is a transmembrane glycoprotein member of the immunoglobulin supergene family of adhesion molecules. CD31 is expressed by stem cells of the hematopoietic system and is primarily used to identify and concentrate these cells for experime... |
| Clonality | Monoclonal |
| Concentration | 0.1 mg/mL |
| Clone Designation | [C31/1395R] |

| | |
|-------------------|---|
| Molecular Weight | ~100 kDa (endothelium) ~130 kDa (platelets) |
| UniProt | P16284 |
| Buffer | PBS, 0.1% rBSA, 0.05% azide |
| Source | Animal |
| Application Notes | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin) 1-2 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris, 1 mM EDTA pH 9.0 for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |