

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

N-Cadherin / Cadherin-2 / CD325 (NCAD) (8C11), CF740 conjugate, 0.1mg/mL, Clone: [8C11], CF 740, Mouse, Monoclonal BOT-BNC741426-100

Article Name	N-Cadherin / Cadherin-2 / CD325 (NCAD) (8C11), CF740 conjugate, 0.1mg/mL, Clone: [8C11], CF 740, Mouse, Monoclonal
Biozol Catalog Number	BOT-BNC741426-100
Supplier Catalog Number	BNC741426-100
Alternative Catalog Number	BOT-BNC741426-100-100UL
Manufacturer	Biotium
Host	Mouse
Category	Antikörper
Application	IHC
Species Reactivity	Human, Mouse
Immunogen	Recombinant human N-cadherin extracellular domain (exact sequence is proprietary)
Conjugation	CF 740
Product Description	This antibody recognizes a protein of ~140 kDa, identified as N-Cadherin (NCAD), also known as CD325. NCAD is a member of the Cadherin superfamily, and consists of five extracellular repeats, a transmembrane domain and a cytoplasmic domain. CD325 def...
Clonality	Monoclonal
Concentration	0.1 mg/mL
Clone Designation	[8C11]

Molecular Weight	130-140 kDa
UniProt	P19022
Buffer	PBS, 0.1% rBSA, 0.05% azide
Source	Animal
Application Notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology (formalin): 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris, 1 mM EDTA pH 9.0 for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user