

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **Factor XIIIa (Coagulation Factor XIIIa Chain) (F13A1/1448), CF740 conjugate, 0.1mg/mL, Clone: [F13A1/1448], CF 740, Mouse, Monoclonal BOT-BNC741448-100**

Article Name	Factor XIIIa (Coagulation Factor XIIIa Chain) (F13A1/1448), CF740 conjugate, 0.1mg/mL, Clone: [F13A1/1448], CF 740, Mouse, Monoclonal
Biozol Catalog Number	BOT-BNC741448-100
Supplier Catalog Number	BNC741448-100
Alternative Catalog Number	BOT-BNC741448-100-100UL
Manufacturer	Biotium
Host	Mouse
Category	Antikörper
Application	IHC, WB
Species Reactivity	Human
Immunogen	Recombinant fragment of human Factor XIIIa protein (aa46-181) (exact sequence is proprietary)
Conjugation	CF 740
Product Description	This antibody recognizes a protein of 83 kDa, which is identified as Factor XIIIa. It has been identified in platelets, megakaryocytes, and fibroblast-like mesenchymal or histiocytic cells in the placenta, uterus, and prostate, monocytes and macropha...
Clonality	Monoclonal
Concentration	0.1 mg/mL

Clone Designation	[F13A1/1448]
Molecular Weight	83 kDa
UniProt	<a href="#">P00488</a>
Buffer	PBS, 0.1% rBSA, 0.05% azide
Source	Animal
Application Notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody ELISA: 2-4 ug/mL for coating order Ab without BSA Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin) 1-2 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user