

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### PGA3 Antibody, Rabbit, Polyclonal NSJ-F40034-0.4ML

|                            |                                                                                                                                                              |
|----------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Article Name               | PGA3 Antibody, Rabbit, Polyclonal                                                                                                                            |
| Biozol Catalog Number      | NSJ-F40034-0.4ML                                                                                                                                             |
| Supplier Catalog Number    | F40034-0.4ML                                                                                                                                                 |
| Alternative Catalog Number | NSJ-F40034-0.4ML                                                                                                                                             |
| Manufacturer               | NSJ Bioreagents                                                                                                                                              |
| Host                       | Rabbit                                                                                                                                                       |
| Category                   | Antikörper                                                                                                                                                   |
| Application                | ELISA, WB                                                                                                                                                    |
| Species Reactivity         | Human                                                                                                                                                        |
| Immunogen                  | A portion of amino acids 45-73 from the human protein was used as the immunogen for this PGA3 antibody.                                                      |
| Product Description        | PGA3 shows particularly broad specificity, although bonds involving phenylalanine and leucine are preferred, many others are also cleaved to some extent.... |
| Clonality                  | Polyclonal                                                                                                                                                   |
| UniProt                    | <a href="#">P0DJD8</a>                                                                                                                                       |
| Purity                     | Antigen affinity                                                                                                                                             |
| Form                       | In 1X PBS, pH 7.4, with 0.09% sodium azide                                                                                                                   |
| Antibody Type              | Primary Antibody                                                                                                                                             |

|                    |                                                                                                                     |
|--------------------|---------------------------------------------------------------------------------------------------------------------|
| Application Dilute | Western blot: 1:1000                                                                                                |
| Application Notes  | Titration of the PGA3 antibody may be required due to differences in protocols and secondary/substrate sensitivity. |