

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### GFAP Antibody, Clone: [ASTRO/789], Mouse, Monoclonal NSJ-V2528-20UG

Article Name	GFAP Antibody, Clone: [ASTRO/789], Mouse, Monoclonal
Biozol Catalog Number	NSJ-V2528-20UG
Supplier Catalog Number	V2528-20UG
Alternative Catalog Number	NSJ-V2528-20UG
Manufacturer	NSJ Bioreagents
Host	Mouse
Category	Antikörper
Application	IHC-P, WB
Species Reactivity	Human, Rat
Immunogen	Full length recombinant protein was used as the immunogen for the GFAP antibody.
Product Description	This mAb recognizes a protein of ~50kDa which is identified as Glial Fibrillary Acidic Protein (GFAP). It shows no cross-reaction with other intermediate filament proteins. GFAP is specifically found in astroglia. GFAP is a very popular marker for lo...
Clonality	Monoclonal
Clone Designation	[ASTRO/789]
UniProt	<a href="#">P14136</a>
Purity	Protein G affinity chromatography

Form	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide
Antibody Type	Primary Antibody
Application Dilute	Western blot: 1-2ug/ml, Immunohistochemistry (FFPE): 0.25-0.5ug/ml for 30 min at RT
Application Notes	Optimal dilution of the GFAP antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 min2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.