

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### CD57 Antibody, Clone: [SPM129], Mouse, Monoclonal NSJ-V2533SAF-100UG

Article Name	CD57 Antibody, Clone: [SPM129], Mouse, Monoclonal
Biozol Catalog Number	NSJ-V2533SAF-100UG
Supplier Catalog Number	V2533SAF-100UG
Alternative Catalog Number	NSJ-V2533SAF-100UG
Manufacturer	NSJ Bioreagents
Host	Mouse
Category	Antikörper
Application	IHC-P
Species Reactivity	Human
Immunogen	Human peripheral blood mononuclear cells were used as the immunogen for the CD57 antibody.
Product Description	Anti-CD57 marks a subset of lymphocytes known as natural killer (NK) cells. Follicular center cell lymphomas often contain many NK cells within the neoplastic follicles. Anti-CD57 also stains neuroendocrine cells and their derived tumors, including c...
Clonality	Monoclonal
Clone Designation	[SPM129]
UniProt	<a href="#">Q9P2W7</a>
Purity	PEG precipitation
Form	1 mg/ml in 1X PBS, BSA free, sodium azide free

Antibody Type	Primary Antibody
Application Dilute	Immunohistochemistry (FFPE): 2-4ug/ml for 30 min at RT
Application Notes	Optimal dilution of the CD57 antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 min2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.