

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

CD57 Antibody Cocktail, Clone: [HNK-1 + NK-1], Mouse, Monoclonal NSJ-V2534IHC-7ML

Article Name	CD57 Antibody Cocktail, Clone: [HNK-1 + NK-1], Mouse, Monoclonal
Biozol Catalog Number	NSJ-V2534IHC-7ML
Supplier Catalog Number	V2534IHC-7ML
Alternative Catalog Number	NSJ-V2534IHC-7ML
Manufacturer	NSJ Bioreagents
Host	Mouse
Category	Antikörper
Application	IHC-P
Species Reactivity	Human
Immunogen	Human peripheral blood mononuclear cells were used as the immunogen for the CD57 antibody cocktail.
Product Description	Anti-CD57 marks a subset of lymphocytes known as natural killer (NK) cells. Follicular center cell lymphomas often contain many NK cells within the neoplastic follicles. Anti-CD57 also stains neuroendocrine cells and their derived tumors, including c...
Clonality	Monoclonal
Clone Designation	[HNK-1 + NK-1]
UniProt	Q9P2W7
Purity	PEG precipitation

Form	Prediluted in 1X PBS, 0.1 mg/ml BSA (US sourced), 0.05% sodium azide, *For IHC use only*
Antibody Type	Primary Antibody
Application Dilute	The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.
Application Notes	Optimal dilution of the CD57 antibody cocktail should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 min2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.