

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

MAGE-1 Antibody, Clone: [MZ2E/838], Mouse, Monoclonal NSJ-V2695SAF-100UG

| | |
|----------------------------|---|
| Article Name | MAGE-1 Antibody, Clone: [MZ2E/838], Mouse, Monoclonal |
| Biozol Catalog Number | NSJ-V2695SAF-100UG |
| Supplier Catalog Number | V2695SAF-100UG |
| Alternative Catalog Number | NSJ-V2695SAF-100UG |
| Manufacturer | NSJ Bioreagents |
| Host | Mouse |
| Category | Antikörper |
| Application | IHC-P |
| Species Reactivity | Human |
| Immunogen | Recombinant human MAGEA1 protein was used as the immunogen for the MAGE-1 antibody. |
| Product Description | Recognizes a protein of 42-46kDa, identified as MAGE-1. This mAb does not cross-react with other members of MAGE-family. Human malignant neoplasms carry rejection antigens that are recognized by the patients autologous, tumor directed and specific, c... |
| Clonality | Monoclonal |
| Clone Designation | [MZ2E/838] |
| UniProt | P43355 |
| Purity | Protein G affinity chromatography |
| Form | 1 mg/ml in 1X PBS, BSA free, sodium azide free |

| | |
|--------------------|---|
| Antibody Type | Primary Antibody |
| Application Dilute | Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT |
| Application Notes | Optimal dilution of the MAGE-1 antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 min.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min. |