

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

CD146 Antibody, Clone: [C146/634], Mouse, Monoclonal NSJ-V2698SAF-100UG

Article Name	CD146 Antibody, Clone: [C146/634], Mouse, Monoclonal
Biozol Catalog Number	NSJ-V2698SAF-100UG
Supplier Catalog Number	V2698SAF-100UG
Alternative Catalog Number	NSJ-V2698SAF-100UG
Manufacturer	NSJ Bioreagents
Host	Mouse
Category	Antikörper
Application	FACS, IF, IHC-P
Species Reactivity	Human
Immunogen	Recombinant human protein was used as the immunogen for the CD146 antibody.
Product Description	The human CD146 gene maps to chromosome 11q23 and encodes a trans-membrane glycoprotein, also designated MCAM, MUC 18, that belongs to the immunoglobulin superfamily and functions as a Ca ²⁺ -independent cell adhesion molecule. CD146 expression is rest...
Clonality	Monoclonal
Clone Designation	[C146/634]
UniProt	P43121
Purity	Protein G affinity chromatography
Form	1 mg/ml in 1X PBS, BSA free, sodium azide free

Antibody Type	Primary Antibody
Application Dilute	Flow cytometry: 0.5-1ug/10 ⁶ cells, Immunofluorescence: 0.5-1ug/ml, Immunohistochemistry (FFPE): 0.5-1ug/ml for 30 min at RT
Application Notes	Optimal dilution of the CD146 antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 minutes2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.