

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### Moesin Antibody, Clone: [MSN/492], Mouse, Monoclonal NSJ-V2718-100UG

Article Name	Moesin Antibody, Clone: [MSN/492], Mouse, Monoclonal
Biozol Catalog Number	NSJ-V2718-100UG
Supplier Catalog Number	V2718-100UG
Alternative Catalog Number	NSJ-V2718-100UG
Manufacturer	NSJ Bioreagents
Host	Mouse
Category	Antikörper
Application	FACS, IF, IHC-P, WB
Species Reactivity	Human
Immunogen	Recombinant full-length human protein was used as the immunogen for the Moesin antibody.
Product Description	Recognizes 78kDa moesin protein. Moesin, a member of the talin-4.1 superfamily, is a linking protein of the sub-membranous actin cytoskeleton. It is expressed in variable amounts in cells of different phenotypes such as macrophages, lymphocytes, fibr...
Clonality	Monoclonal
Clone Designation	[MSN/492]
UniProt	<a href="#">P26038</a>
Purity	Protein G affinity chromatography

Form	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide
Antibody Type	Primary Antibody
Application Dilute	Flow cytometry: 1-2ug/million cells, Immunofluorescence: 1-2ug/ml, Western blot: 1-2ug/ml, Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT
Application Notes	Optimal dilution of the Moesin antibody should be determined by the researcher. 1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min. 2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.