

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Moesin Antibody, Clone: [MSN/492], Mouse, Monoclonal NSJ-V2718IHC-7ML

Article Name	Moesin Antibody, Clone: [MSN/492], Mouse, Monoclonal
Biozol Catalog Number	NSJ-V2718IHC-7ML
Supplier Catalog Number	V2718IHC-7ML
Alternative Catalog Number	NSJ-V2718IHC-7ML
Manufacturer	NSJ Bioreagents
Host	Mouse
Category	Antikörper
Application	IHC-P
Species Reactivity	Human
Immunogen	Recombinant full-length human protein was used as the immunogen for the Moesin antibody.
Product Description	Recognizes 78kDa moesin protein. Moesin, a member of the talin-4.1 superfamily, is a linking protein of the sub-membranous actin cytoskeleton. It is expressed in variable amounts in cells of different phenotypes such as macrophages, lymphocytes, fibr...
Clonality	Monoclonal
Clone Designation	[MSN/492]
UniProt	P26038
Purity	Protein G affinity chromatography

Form	Prediluted in 1X PBS, 0.1 mg/ml BSA (US sourced), 0.05% sodium azide, *For IHC use only*
Antibody Type	Primary Antibody
Application Dilute	The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.
Application Notes	Optimal dilution of the Moesin antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.