

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **c-Myc Antibody, Clone: [9E10.3], Mouse, Monoclonal NSJ-V2744-20UG**

Article Name	c-Myc Antibody, Clone: [9E10.3], Mouse, Monoclonal
Biozol Catalog Number	NSJ-V2744-20UG
Supplier Catalog Number	V2744-20UG
Alternative Catalog Number	NSJ-V2744-20UG
Manufacturer	NSJ Bioreagents
Host	Mouse
Category	Antikörper
Application	IHC-P
Species Reactivity	Human
Immunogen	Amino acids AEEQKLISEEDLLRKRREQLKHKLEQLRNSCA were used as the immunogen for the c-Myc antibody.
Product Description	It recognizes a transcription factor of 64-67kDa, identified as c-myc. Its epitope spans between aa 410-419 (EQKLISEEDL) which is a specific portion of an alpha helical region of human c-myc protein. This mAb shows no cross-reaction with v-myc. c-myc...
Clonality	Monoclonal
Clone Designation	[9E10.3]
UniProt	<a href="#">P01106</a>
Purity	Protein G affinity chromatography

Form	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide
Antibody Type	Primary Antibody
Application Dilute	Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT
Application Notes	Optimal dilution of the c-Myc antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 min2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.