

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Myc Antibody, Clone: [SPM237], Mouse, Monoclonal NSJ-V2745IHC-7ML

Article Name	Myc Antibody, Clone: [SPM237], Mouse, Monoclonal
Biozol Catalog Number	NSJ-V2745IHC-7ML
Supplier Catalog Number	V2745IHC-7ML
Alternative Catalog Number	NSJ-V2745IHC-7ML
Manufacturer	NSJ Bioreagents
Host	Mouse
Category	Antikörper
Application	IHC-P
Species Reactivity	Human
Immunogen	Amino acids AEEQKLISEEDLLRKRREQLKHKLEQLRNSCA from the C-terminus were used as the immunogen for the Myc antibody.
Product Description	Myc, or c-Myc, is involved in the control of cell proliferation and differentiation and is amplified and/or overexpressed in a variety of tumors. Over-expression of c-Myc protein occurs frequently in luminal cells of prostate intraepithelial neoplasia...
Clonality	Monoclonal
Clone Designation	[SPM237]
UniProt	P01106
Purity	Protein G affinity chromatography

Form	Prediluted in 1X PBS, 0.1 mg/ml BSA (US sourced), 0.05% sodium azide, *For IHC use only*
Antibody Type	Primary Antibody
Application Dilute	The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.
Application Notes	Optimal dilution of the Myc antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 min2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.